



Fundy Model Forest

~Partners in Sustainability~

Report Title: Independent Effects of Habitat Amount and Fragmentation on Songbirds in a Forest Mosaic: An Organism-Based Approach

Authors: M. Betts (corresponding author), G.J. Forbes, A.W. Diamond and P.D. Taylor



Natural Resources
Canada

Ressources naturelles
Canada

Canada

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

**INDEPENDENT EFFECTS OF HABITAT AMOUNT AND FRAGMENTATION ON SONGBIRDS IN A
FOREST MOSAIC: AN ORGANISM-BASED APPROACH**

Matthew G. Betts[†] (corresponding author)

Faculty of Forestry and Environmental Management

University of New Brunswick

G.J. Forbes

Faculty of Forestry and Environmental Management

University of New Brunswick

A.W. Diamond,

Atlantic Cooperative Wildlife Ecology Research Network

University of New Brunswick

Phillip D. Taylor

Atlantic Cooperative Wildlife Ecology Research Network

Acadia University

[†] Address for correspondence: 1350 Regent St., Fredericton, New Brunswick, Canada, E3C 2G6, m.betts@unb.ca

1 *Abstract.* The degree to which spatial patterns influence the dynamics and distribution of
2 populations is a central question in ecology. This question is even more pressing in the context
3 of rapid habitat loss and fragmentation which threaten global biodiversity. However, the relative
4 influence of habitat loss and landscape fragmentation – the spatial patterning of remaining
5 habitat – remains unclear. If landscape pattern affects population size, managers may be able to
6 design landscapes that mitigate habitat loss. We present the results of a mensurative experiment
7 designed to test four habitat loss versus fragmentation hypotheses. Unlike previous studies, we
8 measured landscape structure using quantitative, spatially explicit habitat distribution models
9 previously developed for two species: blackburnian warbler (*Dendroica fusca*) and ovenbird
10 (*Seiurus aurocapillus*). We used a stratified sampling design that reduced the confounding of
11 habitat amount and fragmentation variables. Occurrence and reoccurrence of both species were
12 strongly influenced by characteristics at scales greater than the individual territory, indicating
13 little support for the random sample hypothesis. However, the type and spatial extent of
14 landscape influence differed. Both occurrence and reoccurrence of blackburnian warblers were
15 influenced by the amount of poor-quality matrix at 300 and 2000 m spatial extents. The
16 occurrence and reoccurrence of ovenbirds depended on a landscape pattern variable – patch size
17 – but only in cases when patches were isolated. These results support the hypothesis that
18 landscape pattern is important for some species only when the amount of suitable habitat is low.
19 Though theoretical models have predicted such an interaction between landscape fragmentation
20 and composition, to our knowledge this is the first study to report empirical evidence of such
21 non-linear fragmentation effects. Defining landscapes quantitatively from an organism-based
22 perspective may increase power to detect fragmentation effects, particularly in forest mosaics
23 where boundaries between patches and matrix are ambiguous. Our results indicate that

1 manipulating landscape pattern may reduce negative impacts of habitat loss for ovenbird, but not
2 blackburnian warbler. We emphasize that most variance in the occurrence of both species was
3 explained by local scale or landscape composition variables rather than variables reflecting
4 landscape pattern.

5 Key words: Fragmentation, habitat loss, landscape composition, organism-based, patch size,
6 edge effect, matrix, blackburnian warbler, ovenbird.

INTRODUCTION

1
2 The degree to which spatial patterns influence the dynamics and distribution of
3 populations is a central question in ecology (Turner 1989, Levin 1992). This question is even
4 more pressing in the context of rapid habitat loss and fragmentation (FAO 2001, Matthews et al.
5 2001) which are considered principle threats to biodiversity globally (Pimm 1995). It is not
6 surprising that habitat loss has been found to result in population declines (Balmford et al. 2003);
7 behavior (Pulliam and Danielson 1991) and resource availability limit the degree to which
8 increases in population density compensate for reduced habitat area. However, fragmentation,
9 the process of subdividing contiguous habitat into smaller, isolated patches (*sensu* Fahrig 1998),
10 may result in decreased patch colonization and increased rates of local extinction, both of which
11 can result in population declines greater than expected from habitat loss alone (Hanski and
12 Ovaskainen 2000).

13 Determining the relative importance of habitat loss versus fragmentation has been
14 problematic. Because fragmentation occurs through a process of habitat loss in most
15 circumstances, the effects of habitat amount (landscape composition) and fragmentation
16 (landscape configuration) are usually confounded. Simulation modelling studies have addressed
17 this problem by designing artificial landscapes that separate these variables (Fahrig 1998, Hill
18 and Caswell 1999, With and King 1999, Flather and Bevers 2002, Wiegand et al. 2005). In these
19 studies the relative importance of fragmentation appears to depend on the life histories of
20 hypothetical species (Fahrig 1998, With and King 1999). Thresholds have been demonstrated to
21 occur where the influence of fragmentation increases as the amount of suitable habitat decreases
22 in the landscape (Hill and Caswell 1999, Flather and Bevers 2002). However, few empirical
23 studies have been designed to test for the independent effects of landscape composition and

1 configuration (Fahrig 2003). Several of these indicate that habitat amount is the dominating
2 influence on population persistence (McGarigal and McComb 1995, Trzcinski et al. 1999,
3 Heikkinen et al. 2004), but evidence is not unanimous (Villard et al. 1999, Krawchuk and Taylor
4 2003). As in simulation studies, the inconsistency in reported fragmentation effects has been
5 attributed to variability in species' adaptations (e.g., movement ability, home range size,
6 sensitivity to edge; Trzcinski et al. 1999).

7 Variable influences of landscape composition and configuration may also be due to the
8 way in which landscapes have been measured and defined. The theory of island biogeography
9 (MacArthur and Wilson 1967) and metapopulation theory (Levins 1969) form the conceptual
10 underpinnings of landscape ecology, but both are based on simple binary landscapes made up of
11 islands/patches (habitat) and ocean/matrix (non-habitat). While these are easily defined in island
12 archipelagos or simulation models, identifying patches and matrix in real landscapes is often
13 subjective (e.g., Homan et al. 2004, Suorsa et al. 2005) and may not be based in accurate
14 estimates of what a species considers to be habitat (Addicott et al. 1987, Wiens et al. 1993,
15 Wiens 1994). This problem is magnified in forest mosaics where habitat fragments are rarely
16 surrounded by an ecologically neutral or inhospitable environment, and sharpness of edges varies
17 with forest regeneration and succession (Bunnell 1999, Schmiegelow and Mönkönen 2002). The
18 weakness of the patch-oriented approach is that it has failed to capture how individual species
19 perceive and use heterogeneous landscapes (Wiens et al. 1993, Ricketts 2001). Quantitative
20 definitions of suitable habitat at the landscape scale (Boyce and McDonald 1999) offer potential
21 to overcome this weakness. Such spatially explicit habitat models allow more accurate
22 definitions of landscape structure (e.g., patch, matrix, edge) from the perspective of individual
23 organisms (Moilanen and Hanski 1998).

1 The potential influences of landscape composition and configuration on populations can
2 be expressed as four competing hypotheses, each with different implications for how habitat loss
3 and fragmentation affect populations.

4 The *random sample hypothesis* states that small patches are simply random samples of
5 larger patches (Haila 1983). Only factors at the local scale (the spatial extent of individual
6 territories) are important in determining habitat quality. Thus, habitat loss produces a
7 proportional decline in the number of animals living in a particular landscape.

8 In the *landscape composition hypothesis*, the amount of habitat is important, but at larger
9 spatial scales than the individual territory (Fahrig 2003). This distribution pattern could result
10 from preferences of dispersing individuals for breeding sites that are in proximity to conspecifics
11 or heterospecifics (Mönkkönen et al. 1999, Danchin et al. 2004). Large year-to-year
12 environmental variation and short breeding season presumably render such habitat-selection
13 behavior beneficial. Alternatively, movement of organisms could be restricted if there is a large
14 amount of impervious matrix (non-habitat) (Jonsen and Taylor 2000, Goodwin and Fahrig 2002).
15 In a special case of landscape composition hypothesis, abundance of higher quality matrix could
16 decrease the need for habitat at the territory scale. Animals may supplement resources by using
17 the surrounding matrix (Andrén et al 1997, Tubelis et al. 2004).

18 The *landscape fragmentation hypothesis* expects populations to decline linearly with
19 fragmentation, independent of the effects of habitat loss (Villard et al. 1999). Fragmentation
20 effects could result from increased amount of edge, which in some contexts has been found to
21 increase predation (Bátary and Báldi 2004, Marzluff et al. 2004) and reduce food availability
22 (Burke and Nol 1998). Alternatively, fragmentation effects could result from decreases in patch
23 size. Bender et al. (1998) argued that many reported patch size effects could be due simply to

1 increased edge effects in small patches (the geometric effect). Nevertheless, patch size effects
2 could occur independently of edge effect. For species that are reluctant to cross gaps
3 (Desrochers and Hannon 1997), conspecific and heterospecific attraction could result in the
4 selection of larger patches (Connor et al. 2000). Root (1973) argued that densities of organisms
5 might be higher in large patches as a result of increased attraction of moving individuals. Others
6 have predicted that smaller patches should exhibit higher densities due to higher chances of
7 intercepting ground-level dispersers (Bowman et al. 2002), or reduced interspecific competition
8 (MacArthur et al. 1972). Non-threshold landscape configuration effects will result in additive
9 population declines above those that occur as a result of habitat loss.

10 The *non-linear fragmentation hypothesis* states that landscape configuration is important
11 only below some critical amount of suitable habitat (Andrén 1994, Fahrig 1998, Flather and
12 Bevers 2002). Only at low levels of habitat are patches small and isolated enough to result in
13 patch size effects or restrictions in movement (Gardner et al. 1991). This will result in
14 multiplicative (non-linear) effects of fragmentation on habitat loss (i.e., a statistical interaction
15 between landscape configuration and composition).

16 Here, we present the results of a mensurative experiment designed to test these four
17 habitat loss versus fragmentation hypotheses. To do this, we investigated the independent effects
18 of commonly reported composition (amount of suitable habitat, poor-quality matrix) and
19 configuration variables (patch size, edge effect) on the distribution of two species of forest
20 songbirds: blackburnian warbler (*Dendrioca fusca*) and ovenbird (*Seiurus aurocapilla*). Both
21 species are associated with mature forest (Betts et al. *In Press*), are deemed to be area sensitive
22 (Freemark and Collins 1992), and have been in decline in New Brunswick, Canada from 1983-
23 2003 (Sauer et al. 2004) possibly as a result of habitat decline on the breeding grounds (Betts et

1 al. 2003). Unlike previous studies, we adopted a quantitative organism-based approach that relied
2 on independently derived (Betts et al. *In Press*), spatially explicit habitat distribution models to
3 test the effects of landscape composition and configuration.

5 METHODS

6 *Study area*

7 The study was conducted in the Greater Fundy Ecosystem, New Brunswick (NB), Canada
8 (66.08°– 64.96°, 46.08°– 45.47°) (~4000 km²). The Greater Fundy Ecosystem is characterized by
9 89% forest cover, a maritime climate, and rolling topography (elevation 70 – 398m). Forest
10 cover is primarily sugar maple (*Acer saccharum* Marsh.), beech (*Fagus grandifolia* L.), yellow
11 birch (*Betula alleghaniensis* Britt.), red spruce (*Picea rubens* Sarg.), and balsam fir (*Abies*
12 *balsamea* (L.) Mill.). However, black spruce (*P. mariana* Mill.) exists in low-lying areas.
13 Intensive forestry activities (i.e., clearcutting, spruce and pine planting, thinning) have occurred
14 since the early 1970s, resulting in a heterogeneous landscape mosaic where approximately 40%
15 of the study area is mature (>70 years), unmanaged forest (NBDNR 1993).

17 *Spatially explicit habitat models*

18 In a separate study, we developed spatially explicit habitat models for the occurrence of
19 both bird species using local-scale variables derived from GIS as predictor variables (Betts et al.
20 *In Press*) (equations 1 and 2).

$$\begin{aligned} 22 \quad \hat{p} \text{ (blackburnian warbler occurrence)} &= 1 / (\exp (3.58 + 15.63(R) + 1.63(S) \\ 23 \quad + 0.82(Y) - 0.62(M) - 1.42(O) - 0.61(CC) - 0.17(\text{Slope})) + 1) & \quad (1) \end{aligned}$$

$$\hat{p} \text{ (ovenbird occurrence)} = 1 / (\exp (1.50 + 1.79(R) + 1.52(S) + 0.29(Y) - 0.66(M) - 1.29(O) + 0.06(IMW) + 2.86(PINE) + 2.95(SW) + 0.92(TMW) - 0.82(CC)) + 1) \quad (2)$$

Where R, S, Y, M, and O are age class variables representing regenerating, sapling, young, mature, and overmature respectively; CC = crown closure; Slope = slope in degrees; IMW, PINE, SW, and TMW are cover type variables representing shade-intolerant mixedwood, pine, softwood, and shade-tolerant mixedwood, respectively (see Betts et al. *In Press* for age class and cover type details).

GIS land cover data originated from the New Brunswick Forest Inventory (NBDNR 1993), which is based on interpreted and digitized aerial photographs taken in 1993 (1:12,500 scale, colour) and updated to 2000 with satellite imagery (30 m² resolution; Betts et al. 2003). We used Receiver Operating Characteristic curves as a measure of habitat model accuracy. The Area Under the Receiver Operating Characteristic Curve (AUC) describes the relationship between the sensitivity (number of positive observations correctly predicted as positive) and specificity (number of false positive predictions; Hanley and McNeil 1982). The AUC is a single index of classification accuracy that ranges from 0 – 1, and is independent of species prevalence and arbitrary threshold effects (Manel et al. 2001). Models exhibited adequate prediction success (Hosmer and Lemeshow 2000) when tested on independent data from within the Greater Fundy Ecosystem study area (blackburnian warbler AUC = 0.786±0.03 SE, ovenbird AUC = 0.831±0.026 SE) and on independent data from a geographically distinct study area (blackburnian warber AUC = 0.670±0.05 SE, ovenbird AUC = 0.819±0.035 SE; Betts et al. *In Press*).

Landscape variables and study design

Using spatially explicit habitat models, we then developed habitat suitability maps (30 m² resolution) from local-scale GIS models for both blackburnian warbler and ovenbird. These maps were used to identify two landscape configuration variables (patch size, distance from edge) and two landscape composition variables (habitat amount, poor-quality matrix) that are the most likely to influence distribution of forest songbirds (Appendix A; Lichstein et al. 2002a).

To identify habitat patches in the forest mosaic we determined cut points in estimated habitat suitability (\hat{p}) using Receiver Operating Characteristic curves; cut points were identified where the sum of model sensitivity and specificity was maximized (Gu enette and Villard 2005). This approach may overestimate habitat for rarer species, but results in higher prediction success than the use of arbitrary cut points (e.g., $\hat{p}=0.5$; Manel et al. 2001). Cut point values were $\hat{p}=0.41$ and $\hat{p}=0.47$ for blackburnian warbler and ovenbird respectively. As predictive ability of both models was adequate, according to estimates of AUC (see *Spatially explicit habitat models*), we consider these cut points to be reliable. Patch size was measured as the total area of suitable habitat that is separated from other patches by >30 m. Territories of our focal species are unlikely to span gaps of these size (after Villard et al. 1995).

We summed the amount of habitat, weighted by \hat{p} , at two spatial extents for all locations within our study area (300 m, 2000 m radii). These extents reflect those previously found to influence forest passerine habitat use (Drapeau et al. 2000, Mitchell et al. 2001), and likely include the spatial extents relevant to migrant warblers in natal dispersal (Bowman 2003), and extra-territorial movements (Norris and Stutchbury 2001).

We used a randomized stratified sampling design so that samples represented the range of variation in patch size and habitat amount at landscape extents (hereafter “habitat amount”). To

1 serve as a basis for sampling, we defined five patch size categories: 1–20 ha, 21–50 ha, 51–100
2 ha, 101–500 ha, <500 ha, and three habitat amount categories: 0–30%, 31–70%, 71–100%. For
3 the purposes of study design we used the greatest spatial extent (2000 m) to measure habitat
4 amount. We selected sample patches that ensured that all possible combinations of patch size
5 and habitat amount were represented. This involved searching for locations with poorly
6 represented combinations of habitat amount and patch size so that the expected positive
7 correlation between habitat loss and fragmentation was reduced (after Trzcinski et al. 1999). We
8 also selected patches that had the least ambiguous boundaries (i.e., where differences in \hat{p}
9 between within patch habitat, and adjacent non-habitat were greatest). In total, 187 blackburnian
10 warbler and 214 ovenbird patch/landscape combinations were sampled in 2002. Timber
11 harvesting reduced the number of patches to 179 and 203 for blackburnian warbler and ovenbird
12 respectively.

13 In each patch, we established 1–4 sample points beginning 75 m from clearly identifiable
14 forest edges (i.e., roads, recent clearcuts [<10 years]) and proceeding at 250 m intervals toward
15 the patch center. In forest mosaics, such “hard” edges are the most likely to result in decreased
16 habitat quality (Harris and Reed 2002, Manolis et al. 2002). If patches were surrounded by
17 multiple hard edges, as was often the case, location of transect entry point was determined
18 randomly. In 2002, we established 363 individual sample points in blackburnian warbler or
19 ovenbird patches. In 2003, harvesting reduced these to 341.

20 To reflect matrix heterogeneity, we summed the amount of poor-quality matrix at all
21 three spatial extents. We defined poor-quality matrix as areas with very low values of \hat{p} ($<95^{\text{th}}$
22 percentile, $\hat{p}=0.05$). Such poor quality matrix is most likely to be inhospitable for movement
23 (Vega et al. 1998, Haddad and Baum 1999, Belisle and Desrochers 2002). The cut point for poor

1 quality matrix had to be defined arbitrarily since, to our knowledge, no detailed analysis of
2 movement cutpoints in relation to habitat suitability was available in the literature. However, the
3 amount of poor-quality matrix was not highly sensitive to changes in cut point for blackburnian
4 warbler (area $\hat{p}_{0.05} / \hat{p}_{0.1} = 0.793$) or ovenbird (area $\hat{p}_{0.05} / \hat{p}_{0.1} = 0.843$).

5 One drawback to a patch-based research design can be that landscape scale sample size is
6 very low (because multiple patches exist within a single landscape), limiting the ability to make
7 landscape-scale inferences (Fahrig 2003). We avoided this problem by characterizing the
8 landscape surrounding each individual sample point and then accounting for lack of
9 independence of points within a patch by using mixed models. The independence of the resulting
10 landscapes were also tested using spatial autocorrelation (see *Statistical analysis*). A key
11 advantage of this patch-based design was that it allowed us to precisely characterize, separate,
12 and test the independent effects of fragmentation and habitat amount. Once patches are
13 identified, patch size is an unambiguous and easily measured variable whereas landscape-scale
14 patch metrics (e.g., mean patch size, patch size coefficient of variation) are both difficult to
15 interpret (Gustafson 1998) and control for in study designs.

16 *Bird sampling*

17 We conducted fixed-radius point counts of forest passerines (Ralph et al. 1995) at each
18 sample point within the period 03 June–11 July in both 2002 and 2003. Three counts of 5-
19 minute duration were conducted on separate occasions between 05:30–11:00 AST. All male
20 birds seen or heard during this time period within a 50 m radius were recorded as ‘present’.
21 Birds flying overhead were not used in data analysis. Because mean bird counts per station
22 tended to be low (<2) for both species, and because we were interested in estimating probability
23 of occurrence, we reduced relative abundance data to presence/absence for use in binomial

1 models. Because the number of times a bird is observed at a location in successive years may be
2 an indicator of habitat quality (Hames et al. 2001, Rodenhouse et al. 2003), we also identified
3 sites that were occupied by each species in both 2002 and 2003 (hereafter “reoccurrence”).

4 *Vegetation sampling*

5 Evaluating the hypothesis that vegetation at the local scale alone can explain variation in
6 forest bird occurrence (random sample hypothesis) required detailed information about
7 vegetation composition and structure at local scales. At each point count location we counted
8 and identified to species all woody stems >2cm diameter at breast height (dbh) within a 20 x 10
9 m plot (0.02 ha, 2.5% of 50 m point count circle; Bowman et al. 2001). To reduce variables used
10 in analysis, we collapsed tree data into dbh categories: 10–30 cm dbh and >30cm dbh, and two
11 species groups (coniferous, deciduous; Appendix A). We calculated basal area using all trees >2
12 cm dbh. Shrubs (woody stems < 2 cm diameter and > 0.5 m tall) were tallied in a 20 x 2 m plot
13 nested within the larger plot. Canopy cover was estimated with a vertical viewing tube 10 cm
14 long and 3 cm inside diameter and fitted with crosshairs. Readings were taken by counting the
15 number of times crosshairs intersected with canopy foliage at 2 m intervals around the perimeter
16 of the 20 x 10 m plot (Emlen 1967). We estimated the number of mature spruce (>20 cm dbh; a
17 known requirement for blackburnian warbler nesting; Morse 1994, Young et al. 2005) within 50
18 m of each sample point. The number of spruce was recorded as one of the following density
19 classes: 0, 1–5, 6–10, 11–50, 51–75, 76–100, >100 stems/ 50m radius point count plot (after
20 Young et al. 2005).

21 For analysis, we selected habitat variables based on local scale habitat relationships
22 observed in the study region (Betts et al. *In Press*) and elsewhere (for reviews see Morse 1994,
23 Van Horne and Donovan 1994; Appendix A).

1 *Statistical Models*

2 Our study design made it necessary to account for the potential lack of independence
3 among multiple points nested within a single patch. We applied generalized linear mixed models
4 (GLMMs) using Penalized Quasi-Likelihood to determine parameter estimates. These models
5 penalize estimated standard errors for potential lack of independence due to grouping, and are
6 thus appropriate for modeling the dependence among outcome variables inherent in clustered
7 data (see Breslow and Clayton 1993 for details). Individual patches were treated as random
8 effects and all other variables as fixed effects. All models were fit in R 2.0.1 (R Development
9 Core Team 2004) statistical program using the GLMM routine with a binomial family (Bates and
10 Sarkar 2005).

11 We used the information-theoretic approach as a model selection procedure (Burnham
12 and Anderson 2002). The advantage of this approach is that it allows one to measure and reflect
13 model selection uncertainty. Models with lower Akaike's Information Criterion (AIC) values are
14 better fitting and more importantly, the relative likelihood of each model in relation to the best
15 model can be determined using evidence ratios derived from AIC values. The evidence ratio can
16 be interpreted as the number of times less likely model i is than the model with the lowest AIC.
17 In cases where model selection uncertainty existed, we used AIC weights to determine the
18 relative importance of models. Weights were summed over the subset of models that included
19 variable x_1 . AIC weights can be interpreted as the Bayesian posterior probabilities for the model
20 set, describing their relative likelihoods of best fitting the data (Zabel et al. 2003).

21 To test for the independent effects of landscape composition and configuration on the
22 occurrence of both forest bird species, we used a sequential model-building approach. First, we
23 fit models for occurrence and reoccurrence, using only local habitat variables. If two or more

1 predictor variables were highly correlated ($r > 0.7$), we included variables that fit the data better
2 (from visual inspections of residual plots and explained deviance). Because little model selection
3 uncertainty existed among local habitat models for either species, we considered the best models
4 to be the ones with the smallest AIC values.

5 Once best local models had been determined, we used AIC to assess the weight of
6 evidence for, or against, four habitat loss versus fragmentation hypotheses. We applied the
7 following model building approach: (1) To control for local-scale variability, best local habitat
8 variables were always retained (Lichstein et al 2002b). (2) To avoid multicollinearity, we did not
9 include any landscape habitat and poor-quality matrix variables from the same spatial extent in
10 the same models. (3) Even though we explicitly designed our sampling to separate the
11 confounding of configuration and cover, some degree of correlation occurred among landscape
12 variables (Table 1). To consider the effects of landscape configuration over and above the
13 influence of habitat amount, we included at least one large-extent habitat amount variable (2000
14 m) in addition to any configuration terms (Fahrig 2003, Krawchuk and Taylor 2003). (4) We
15 evaluated the non-linear fragmentation hypothesis by examining interactions between both
16 configuration variables (patch size, edge) and composition variables (habitat, poor-quality matrix
17 amounts at a 2000 m extent) (Appendix A). Support for such an interaction would indicate a
18 non-linear relationship between landscape composition and configuration and support for the
19 non-linear fragmentation hypothesis (Tzrcinski et al. 1999). (5) If models with one or two
20 landscape variables ranked within 4 AIC units of the best model, these were combined to
21 determine if model fit could be improved (Zabel et al. 2003). These model-building rules
22 resulted in a candidate set of 21–22 models for each species in each year and for reoccurrence.
23 We did not detect overdispersion in either local or landscape GLMM models ($\hat{c} < 1$).

1 Collinearity in explanatory variables often hampers the detection of the independent
2 effects of environmental variables. We used a variance partitioning approach (Chevan and
3 Sutherland 1991, Borcard et al. 1992, MacNally 2000) to determine the independent
4 contributions of local habitat, landscape composition, and landscape configuration variables to
5 explained variance. Partial explained deviance values for local, landscape composition, and
6 landscape configuration variables were calculated as the increase in explained deviance in
7 logistic regression models due to the inclusion of variables from each predictor set after
8 controlling for variables from all other sets (Venables and Ripley 2001).

9 The presence of spatial autocorrelation is not simply a statistical problem to be avoided –
10 it can provide important information about the ecology of species (Legendre 1993). We used
11 correlograms of Moran's I (hereafter I) to test for autocorrelation in Pearson residuals of all
12 regression model sets (Kaluzny et al. 1996; Klute et al. 2002). We standardized I by dividing by
13 its maximum value (after Haining 1990; Lichstein et al. 2002b). Because the shortest distance
14 between sample points in both study areas was ~250–350 m, our lag intervals were at 350 m up
15 to a maximum distance of 7000 m. We used randomization tests (999 permutations) to
16 determine the probability of observing a value of I as large as the observed value. For each
17 correlogram, the significance of I for each lag distance was calculated using a Bonferroni
18 correction for multiple tests (after Lichstein et al. 2002b). To test the hypothesis that landscape
19 composition effects may be due to aggregation (Lichstein et al. 2002b), we tested for spatial
20 autocorrelation in residuals of bird habitat models that included only local-scale vegetation.
21 Because GLMMs effectively removed all spatial dependency of points in proximity within the
22 same patch (see *Results*), we used fixed effects models (GLMs) in these tests for spatial

1 autocorrelation. We predicted that local models should exhibit spatial autocorrelation, but that
2 the inclusion of landscape terms should account for this aggregative effect.

3 **RESULTS**

4 *Blackburnian warbler*

5 Blackburnian warbler occurrence and reoccurrence at the local scale tended to be
6 associated with large (>30 cm dbh) tree density, mixed coniferous-deciduous forest, and a high
7 percentage of canopy cover (Table 2). However, we found little support for the random sample
8 hypothesis; the model including only local variables tended to perform poorly and was 167.7
9 (Δ AIC 10.23), 11.9 (Δ AIC 4.95) and 40.0 (Δ AIC 7.37) times less likely to fit the data than the
10 top-ranked model in 2002, 2003 and for reoccurrence respectively (Table 3). Evidence from AIC
11 and variance partitioning supported the landscape composition hypothesis. Occurrence of
12 blackburnian warbler in both years and reoccurrence was best predicted by models containing
13 only landscape composition variables (2002: top 4 models $\sum w_i = 0.753$; 2003: top 3 models $\sum w_i$
14 = 0.692; reoccurrence: top 5 models $\sum w_i = 0.745$; Appendices B-D). The contribution of
15 landscape variables to independently explained variance in blackburnian warbler occurrence was
16 25% (2002), 22% (2003) and 29% (reoccurrence; Fig. 1A). Models containing fragmentation
17 variables tended to have only weak support and explained little variance (Table 3, Fig. 1). Thus,
18 in this species, there was little support for either the fragmentation or non-linear fragmentation
19 hypothesis.

20 We found considerable uncertainty about the most important spatial extent for
21 blackburnian warbler. Occurrence in 2002 and reoccurrence were best predicted by the amount
22 of poor-quality matrix at both 300 m and 2000 m spatial extents (Table 2, Appendices B and D.
23 Controlling for local-scale variation, the species was less likely to occur and reoccur in

1 landscapes with large amounts of matrix at these extents. In 2003, the best model indicated that
2 the occurrence of blackburnian warbler was most reliably predicted only by the amount of matrix
3 within 300 m (Table 2, Appendix C).

4 *Ovenbird*

5 Ovenbird occurrence and reoccurrence at the local scale was positively correlated with
6 canopy cover, basal area of deciduous trees and leaf litter (Table 4). However, as with
7 blackburnian warbler, we found little support for the random sample hypothesis; the ovenbird
8 model including only local variables was not supported in 2002 (evidence ratio [ER]: 28 991.3,
9 Δ AIC 20.55), or for reoccurrence (ER: 30.7, Δ AIC 6.85; Table 5), but had weak support in 2003
10 (ER: 6.7, Δ AIC 3.79). The proportion of variance independently explained by landscape
11 variables was 62%, 63% and 65% for 2002, 2003, and reoccurrence respectively (Fig. 1B).

12 We found considerable support for the non-linear fragmentation hypothesis for ovenbird.
13 The occurrence of this species in 2002 and reoccurrence was best predicted by both composition
14 and configuration variables (Table 5, Appendices E and G); AIC weights of models containing
15 interactions between landscape configuration and composition constituted the best two models in
16 2002 ($\sum w_i = 0.973$) and the best three models for reoccurrence ($\sum w_i = 0.881$; Appendices E and
17 G). However, in 2003, models supporting the landscape composition hypothesis were reasonable
18 competitors with the composition-configuration interaction models (interaction models $\sum w_i =$
19 0.414, composition models $\sum w_i = 0.358$; Appendix F). We found very little support for the
20 fragmentation hypothesis; models containing configuration variables in the absence of
21 composition-configuration interaction terms were not supported (Table 5).

22 Patch size, rather than edge effect was the most important configuration variable in
23 ovenbird models; edge effect appeared in none of the top models (Δ AIC<4) in 2002 or for

1 reoccurrence and only one of the top models in 2003 ($\Delta AIC=3.7$; Appendices E-G). Ovenbird
2 was less likely to occur in small patches, but only when those patches were isolated (i.e., in
3 landscapes containing relatively small amounts of habitat at the 2000 m spatial extent; Fig. 2). In
4 2002, ovenbird occurrence was positively correlated with the amount of habitat at both 300 m
5 and 2000 m spatial extents (Table 4). However, there was variation in the relative importance of
6 these extents between years and for reoccurrence (Table 4, Appendices E-G).

7 *Spatial autocorrelation*

8 We did not detect spatial autocorrelation in the residuals of either local or landscape
9 GLMMs for blackburnian warbler or ovenbird. Nor did we detect spatial autocorrelation in
10 ovenbird GLMs including landscape variables; residuals showed little spatial pattern, indicating
11 that the assumption of independent errors was not violated (Lichstein et al. 2002b). However,
12 residuals of local-scale fixed effects models (GLMs) were spatially autocorrelated for ovenbird
13 in 2002 (350 m: $I=0.16$, $P=0.01$), 2003 (350 m: $I=0.12$, $P=0.04$; 700 m: $I=0.18$, $P=0.003$) and
14 reoccurrence (700 m: $I=0.13$, $P=0.02$) and for blackburnian warbler in 2002 (700 m: $I=0.17$,
15 $P=0.001$, 1400 m: $I=0.19$, $P=0.002$). Top-ranked fixed effects models that included landscape
16 variables exhibited spatial autocorrelation in residuals only in 2002 for blackburnian. In this
17 case, autocorrelation existed to a lesser degree than in the local-scale model and at only one
18 distance (700 m: $I=0.15$, $P=0.01$).

19 **DISCUSSION**

20 *Landscape effects and the random sample hypothesis*

21 Our results provide little support for the random sample hypothesis. Occurrence and re-
22 occurrence of both species depended on forest characteristics at spatial extents greater than the
23 individual territory. This suggests that forest landscape data do not consistently match the

1 random sample hypothesis (*contra* Mönkkönen and Reunanen 1999). More recent studies
2 conducted in forest mosaics have found that variables at landscape extents are significant
3 predictors of bird occurrence. However, the importance of landscape variables has tended to be
4 substantially less than local variables (Norton et al. 2000, Hagan and Meehan 2002). For
5 instance, Lichstein et al. (2002a), one of the few forest mosaic studies to use a variance
6 partitioning approach, found that landscape variables accounted for 0-24% of independently
7 explained variance. Our results for blackburnian warbler are similar; the amount of variance
8 explained by landscape variables was low in comparison to local scale variables (22-29%).
9 However, over 60% of the independently explained variance in ovenbird occurrence was due to
10 landscape variables. Comparatively strong landscape effects in our study are likely due to three
11 non-mutually exclusive factors.

12 First, our samples were located in a subset of the entire forest where previously derived
13 local-scale statistical models predicted the occurrence of both species (Betts et al. *In Press*).
14 Several previous studies have commonly sampled all broadly-defined cover types within a high
15 contrast forest (e.g., Drapeau et al. 2000, Lichstein 2002a). Because nesting and foraging
16 requirements of many bird species are relatively stereotyped (Holmes and Sherry 1986), it is not
17 surprising that landscape variables have been found to be comparatively less important; if
18 appropriate nesting and foraging substrate are not available, a species is unlikely to occur
19 regardless of how much appropriate habitat there is at broader spatial extents. However, in the
20 current study, we held local level variation constant through a combination of study design and
21 statistical control, which provided more power to detect landscape effects.

22 Second, detection of only minor landscape effects may result from a relatively high
23 proportion of suitable habitat in a study landscape (Tewksbury et al. 1998, Norton et al. 2000,

1 Lichstein et al. 2002a). In a simulation study, Andrén (1996) found that power to reject the
2 random sample hypothesis is lower in landscapes with high proportions of suitable habitat. In
3 our study, the range in the proportion of suitable habitat at the largest spatial extent was broad for
4 both species (blackburnian warbler $\bar{x}=28.5\pm 5.0\%$, range=9.2-57.1%; ovenbird $\bar{x}=37.9\pm 0.7\%$,
5 range=11.6-74.7%; percentages were calculated using amount of highest quality habitat [$\hat{p}=1.0$]
6 within 2000 m as a denominator). These values encompass the range of values (Andrén 1996)
7 that would allow detection of landscape effects for species that are either area sensitive or poor
8 dispersers.

9 Third, quantitative, organism-based approaches to defining landscape characteristics are
10 still rare (Reunanen et al. 2002). Such an approach is particularly important in a forest mosaic
11 where distinctions between habitat and matrix are less discrete than in agricultural mosaics or
12 island archipelagos (Mönkkönen and Reunanen 1999). Previous studies have relied on arbitrary,
13 or at least general definitions of suitable habitat at landscape extents (Trzcinski et al. 1999,
14 Fischer et al. 2004).

15 *Landscape composition versus fragmentation hypotheses*

16 Through a combination of study design and statistical methods, we separated the often-
17 confounded effects of landscape configuration and composition. The two species we examined
18 responded to landscape structure differently. After controlling for local habitat and landscape
19 composition, blackburnian warbler models containing configuration variables had very little
20 support; our results for this species support the landscape composition hypothesis. Hagan and
21 Meehan (2002) and MacFaden and Capen (2002) found weak, but significant landscape
22 composition effects on this species. The lack of configuration effects is consistent with most

1 studies that have attempted to separate the effects of habitat loss from fragmentation *per se* (see
2 Fahrig 2003 for review).

3 In contrast, ovenbird distribution in 2002 and reoccurrence were strongly influenced by
4 landscape pattern. We found a positive influence of patch size on ovenbird occurrence, but only
5 when the amount of suitable habitat in the landscape was low. This result supports, for the first
6 time to our knowledge, the non-linear fragmentation hypothesis (Fahrig 2003). In forest-
7 agricultural landscapes, Villard et al (1999) found this species to be positively correlated with the
8 amount of forest cover at the 2.5 x 2.5 km extent, but not with fragment area or edge effect.
9 Several studies have reported greater likelihood of ovenbird occurrence in contiguous forest than
10 in small, isolated patches (Hannon and Schmiegelow 2002, Nol et al. 2005). Lee et al. (2002)
11 found that forest cover explained the most variation in ovenbird abundance, but that ovenbird
12 density was lower in large patches. Lee et al. (2002) speculated that this negative patch size
13 effect was due to habitat supplementation from foraging outside of small patches. However,
14 none of these studies tested for interactions between patch size and habitat amount (a non-linear
15 effect).

16 Numerous theoretical studies have found non-linear responses by species to
17 fragmentation (Fahrig 1998, With and King 1999, Flather and Bevers 2002, Wiegand et al.
18 2005). Thresholds have been predicted to occur for species with low vagility, non-ephemeral
19 habitat, high site-fidelity, and high mortality in non-breeding habitat areas (Fahrig 1998). Given
20 this narrow range of conditions, and the logistical difficulty of separating the confounding effects
21 of landscape composition and configuration, it is perhaps not surprising that few empirical
22 studies have reported evidence for non-linear fragmentation effects. Ovenbirds are site faithful
23 (Van Horne and Donovan 1994) and establish territories in deciduous forest, a forest type that is

1 relatively stable (Lorimer 1977). However, dispersal distances of juveniles, and mortality in the
2 matrix are unknown (Villard et al. 1995). More information on these life history characteristics
3 is required before the conditions of fragmentation sensitivity put forward by theoretical models
4 can be validated.

5 We found almost no support for pure fragmentation effects, uninfluenced by the amount
6 of suitable habitat, for either species. This is consistent with numerous studies (Sallabanks et al.
7 2000, Drapeau et al. 2000, MacFaden and Capen 2002, but see Villard et al. 1999). The lack of
8 support for the fragmentation hypothesis has been anticipated by a number of researchers since
9 the initial stages of habitat loss, patches remain relatively well connected (Turner et al. 1989,
10 Gardner et al. 1991). Thus, if the amount of suitable habitat in landscape is high, species are
11 unlikely to respond to, or even perceive, gaps between patches.

12 *Inferring process from pattern*

13 In simulation studies, metrics summarizing habitat amount, or non-habitat in a landscape-
14 extent circle surrounding a patch were the best measures of patch isolation (Bender et al. 2003,
15 Tischendorf et al. 2003). For ovenbird, we found that amount of habitat at the 300 m and 2000 m
16 extents were important predictors of occurrence and reoccurrence. Small patches in landscapes
17 with low amounts of suitable habitat were less likely to be occupied. In landscapes with high
18 proportions of suitable habitat, ovenbird movement is unlikely to be restricted by the small gaps
19 that occur. However, as the proportion of suitable habitat declines, gaps between patches
20 become greater and emerge as potential barriers to moving adults and juveniles (Hinsley 2000).
21 Indeed, previous research in a forest mosaic indicates that this species may avoid crossing gaps
22 during the breeding season (Robichaud et al. 2002, Bayne et al. 2005) and is less likely to move
23 through landscapes with low forest cover (Belisle et al. 2001, Gobeil and Villard 2002).

1 Ovenbirds are ground nesters and foragers that rely on invertebrates found in deciduous litter
2 (Burke and Nol 1998). This specialization may require dispersal or extra-territorial movements
3 through a narrow range of forest types; if the species moves preferentially through what we
4 defined as ovenbird habitat, connected patches will be used more frequently in landscapes with
5 low proportions of suitable habitat.

6 Our spatial autocorrelation analysis indicates that responses to landscape structure by this
7 species may be driven by a tendency to aggregate. Even after controlling for the effects of local
8 vegetation, ovenbird tended to be positively autocorrelated at 350 m (occurrence 2002, 2003) and
9 700 m scales (occurrence 2002, 2003, reoccurrence); clustering seems to occur independent of
10 local habitat structure. Models including landscape terms were not autocorrelated. Landscape
11 pattern thus seems to account for the observed aggregation in this species. Forest birds have
12 been shown to rely on cues from conspecifics (Danchin et al. 2004) or heterospecifics
13 (Mönkkönen et al. 1999) for information about habitat quality. Small, isolated patches would be
14 less likely to contain large numbers of cue-providing individuals, reducing the opportunity to
15 capitalize on social information (Danchin et al. 2004) and thus decreasing settlement rates.
16 Isolation effects could also result from reduced opportunities for extra-pair fertilization in birds—
17 an occurrence that is potentially much more common than previously assumed (Chuang-Dobbs et
18 al. 2001, Webster et al. 2001) and may boost productivity (Holmes et al. 1992). Presumably, it is
19 more risky to foray into patches at greater distances, as it requires more energy and longer
20 periods away from defended territories (Woolfenden et al. 2005).

21 For blackburnian warbler, the variables representing the amount of poor-quality matrix at
22 both spatial extents were the most important predictors. Forest types that are clearly non-habitat
23 ($\hat{p} < 0.05$) may constitute barriers to this species (Belisle et al. 2001). Avoidance of landscapes

1 with poor-quality matrix could reflect either supplementation behaviour by this species, or
2 restricted movement of juveniles. This result for blackburnian warbler supports previous
3 research indicating that the landscape matrix quality can influence species movement and
4 distributions (Lindenmayer et al. 2002, Brotons et al. 2003, Baum et al. 2004, Tubelis et al.
5 2004).

6 Why did we detect a patch size effect for ovenbird but not blackburnian warbler?

7 Because our initial habitat model for blackburnian warbler had lower prediction success than the
8 ovenbird model, it is possible that we did not define patches as effectively. However, we
9 situated sample points in patches where the distinction between habitat and non-habitat was the
10 least ambiguous (see *Methods*). This reduced the likelihood that areas beyond defined patch
11 boundaries were still adequate habitat. Alternatively, differences may be due to foraging breadths
12 of these species. Blackburnian warblers have specialized nesting requirements, but a broader
13 foraging niche than ovenbirds (Morse 1994). This may allow movement through a wider range of
14 forest types, resulting in a more permeable matrix (McComb 1999). For instance, Vega et al.
15 (1998) found that juvenile wood thrushes (*Hylocichla mustelina*) tended to disperse where food
16 availability was high, rather than remaining solely in mature deciduous forest that characterizes
17 natal territories.

18 *Conservation implications*

19 Our most important management-related finding is that habitat loss is unlikely to result in
20 a proportional decline in the number of animals. For both species, landscape composition is an
21 important predictor of occurrence and reoccurrence. For ovenbird, landscape configuration is
22 also important. If timber harvesting creates high amounts of low quality matrix for blackburnian
23 warblers, population decline will occur more rapidly than expected from simple loss of habitat.

1 For this species, enhancing the permeability of the landscape matrix may increase occurrence in
2 sites with appropriate local scale habitat.

3 Our finding that the probability of ovenbird occurrence is greater in larger patches, but
4 only in landscapes with low proportions of suitable habitat, suggests that manipulating landscape
5 pattern (e.g., leaving some large patches unharvested) may mitigate the negative effects of
6 habitat loss for some species. This is particularly important in light of current declines in mature
7 forest in New Brunswick due to timber harvesting (Betts et al. 2003). However, altering the
8 spatial configuration of blackburnian warbler habitat is unlikely to have beneficial results.
9 Further, non-linear fragmentation effects are probably context specific; it is possible that as the
10 amount of suitable habitat at a broader regional scale declines, interactions between landscape
11 configuration and composition could shift. This hypothesis may be tested through simulation
12 (e.g., Donovan and Thompson 2001, Larson et al. 2004); it is usually untenable to manipulate
13 habitat amount experimentally at such large scales. However, an organism-based approach
14 applied to multiple species may offer the opportunity to test this hypothesis empirically since
15 each species simultaneously perceives the region as containing different proportions of suitable
16 habitat. Species with similar degrees of habitat specialization to ovenbird should exhibit greater
17 responses to landscape pattern if there is less suitable habitat at the regional scale.

18 That the majority of variance in the occurrence of both species was explained by
19 composition variables at the local and landscape scale gives support to the basic premise that
20 maintaining the amount of habitat in the landscape is critical. However, our observation of a
21 non-linear fragmentation effect for ovenbird further suggests that managers can justifiably
22 pay attention to landscape configuration effects to further enhance the conservation value of
23 landscapes for some species.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Acknowledgments

We are grateful to A. Breton, S. Makepeace, J. Nocera, N. Simon, two anonymous reviewers, and J. Marzluff who provided useful comments on this manuscript. Also, thanks are due to J. Cormier, A. Frank, G. George, A. Hadley, R. Kuzyk, D. Mitchell, K. Pistak, A. Whaley, E. Widden, and B. Zitske who assisted in the field component of this study, and to S. White and R. Taylor who assisted in compilation of GIS data. Financial support was provided by the New Brunswick Environmental Trust Fund, New Brunswick Department of Natural Resources, the Fundy Model Forest, Fundy National Park, and a doctoral fellowship to MGB from the Natural Sciences and Engineering Research Council (NSERC). This is ACWERN publication No. UNB-49.

Literature Cited

Addicott, J.F., J.M. Aho, M.F. Antolin, D.P. Padilla, J.S. Richardson and D.A. Soluk. 1987. Ecological neighbourhoods: scaling environmental patterns. *Oikos* **49**:340-346.

Andrén H. 1994. Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos* **71**:355-366.

Andrén H. 1996. Population responses to habitat fragmentation: statistical power and the random sample hypothesis. *Oikos* **76**:235-242.

Andrén, H., A. Delin and A. Seiler. 1997. Population response to landscape changes depends on specialization to different landscape elements. *Oikos* **80**:193-195.

Balmford, A., R.E. Green, and M. Jenkins. 2003. Measuring the changing state of nature. *Trends in Ecology and Evolution* **18**:326-330.

- 1 Batáry, P. and A. Báldi. 2004. Evidence of an edge effect on avian nest success. *Conservation*
2 *Biology* **18**:389-400.
- 3 Bates, D. and D. Sarkar. 2005. Linear mixed effects models using S4 models. LME 4 package for
4 R.
- 5 Baum, K., K.J. Haynes, F.P. DelleMuth, and J.T. Cronin. 2004. The matrix enhances the
6 effectiveness of corridors and stepping stones. *Ecology* **85**:2671-2676.
- 7 Bayne, E.M., S.L. Van Wilgenburg, S. Boutin, and K.A. Hobson. 2005. Modeling and field-
8 testing of Ovenbird (*Seiurus aurocapillus*) responses to boreal forest dissection by energy
9 sector development at multiple scales. *Landscape Ecology* **20**:203-216.
- 10 Belislé, M., A. Desrochers, and M.-J. Fortin. 2001. Influence of forest cover on the movement of
11 forest birds: a homing experiment. *Ecology* **82**:1893-1904.
- 12 Belislé, M. and A. Desrochers. 2002. Gap-crossing decisions by forest birds: an empirical basis
13 for parameterizing spatially-explicit, individual-based models. *Landscape Ecology* **17**:
14 219-231.
- 15 Bender D.J., T.A. Contreras and L. Fahrig. 1998. Habitat loss and population decline: a meta-
16 analysis of the patch size effect. *Ecology* **79**:517-533.
- 17 Bender, D.J., L. Tischendorf, and L. Fahrig. 2003. Using patch isolation metrics to predict animal
18 movement in binary landscapes. *Landscape Ecology* **18**:17-39.
- 19 Betts, M.G., A.W. Diamond, G.J. Forbes, M.-A. Villard, and J. Gunn. 2005. The importance of
20 spatial autocorrelation, extent and resolution in predicting forest bird occurrence.
21 Accepted to: *Ecological Modelling*, *in press*.
- 22 Betts, M.G., S.E. Franklin and R.S. Taylor. 2003. Interpretation of landscape pattern and habitat
23 change for local indicator species using satellite imagery and geographic information

- 1 system data in New Brunswick, Canada. *Canadian Journal of Forest Research* **33**:1821-
2 1831.
- 3 Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the spatial component of
4 ecological variation. *Ecology* **73**:1045-1055.
- 5 Bowman, J., N. Cappuccino, and L. Fahrig. 2002. Patch size and population density: the effect of
6 immigration behavior. *Conservation Ecology* 6:9. <http://www.consecol.org/vol6/iss1/art9>.
- 7 Bowman, J. 2003. Is dispersal distance of birds proportional to territory size? *Canadian Journal*
8 *of Zoology* **81**:195-202
- 9 Boyce, M.S. and L.L. MacDonald. 1999. Relating populations to habitats using resource
10 selection functions *Trends in Ecology and Evolution* **14**:268-272.
- 11 Breslow, N.E., and D.G. Clayton. 1993. Approximate inference in Generalized Linear Mixed
12 Models. *Journal of the American Statistical Association* **88**:9-24.
- 13 Brotons, L., M. Mönkkönen, and J.L. Martin. 2003. Are fragments islands? Landscape context
14 and density-area relationships in boreal forest birds. *American Naturalist* **162**: 343-357.
- 15 Bunnell, F.L. 1999. What is a habitat island? Pages 1-31 *in* J.A. Rochelle, L.A. Lehmann, and J.
16 Wisniewski, editors. *Forest Fragmentation : wildlife and management implications*. Brill
17 Academic Publishing, Leiden, The Netherlands.
- 18 Burke D.M. and E. Nol. 1998. Influence of food abundance, nest-site habitat, and forest
19 fragmentation on breeding ovenbirds. *Auk* **115**:96-104.
- 20 Burnham, K.P. and D.R. Anderson. 2002. *Model Selection and Multimodel Inference: A*
21 *Practical Information-Theoretic Approach*. Springer-Verlag, New York.
- 22 Chevan, A, and M. Sutherland. 1991. Hierarchical partitioning. *American Statistician* **45**:90-96.

- 1 Chuang-Dobbs, H.C., M.S. Webster, and R.T. Holmes. 2001. Paternity and parental care in the
2 black-throated blue warbler, *Dendroica caerulescens*. *Animal Behaviour* **62**:83-92.
- 3 Connor, E.F., A.C. Courtney and J.M. Yoder. 2000. Individuals-area relationships: the
4 relationship between animal population density and area. *Ecology* **81**:734-748.
- 5 Danchin, E., L.-A. Giraldeau, T.J. Valone, and R.H. Wagner. 2004. Public information: from
6 nosy neighbors to cultural evolution. *Science* **305**:487-491.
- 7 Desrochers A. and S.J. Hannon 1997. Gap crossing decisions by forest songbirds during the post-
8 fledging period. *Conservation Biology* **11**:1204-1210.
- 9 Donovan, T.M., and F.R. Thompson. 2001. Modeling the ecological trap hypothesis: a habitat
10 and demographic analysis for migrant songbirds. *Ecological Applications* **11**:871-882.
- 11 Drapeau, P., A. Leduc, J.F. Giroux, J.P. Savard, , Y. Bergeron, and W.L. Vickery. 2000.
12 Landscape-scale disturbances and changes in bird communities of boreal mixed-wood
13 forests. *Ecological Monographs* **70**:423-444.
- 14 Emlen, J.T. 1967. A rapid method for measuring arboreal canopy cover. *Ecology* **48**:158-160.
- 15 Fahrig, L. 1998. When does fragmentation of breeding habitat affect population survival?
16 *Ecological Modelling* **105**:273-292.
- 17 Fahrig, L. 2003. Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology*
18 *Evolution and Systematics* **34**:487-515.
- 19 Fischer, J., D.B. Lindenmayer, I. Fazey. 2004. Appreciating ecological complexity: habitat
20 contours as a conceptual landscape model. *Conservation Biology* **18**:1245-1253.
- 21 Flather, C.H. and M. Bevers. 2002. Patchy reaction-diffusion and population abundance: the
22 relative importance of habitat amount and arrangement. *American Naturalist* **159**:40-56.

- 1 Food and Agriculture Organization. 2001. Global Forest Resources Assessment 2000: Main
2 Report, FAO.
- 3 Freemark, K.E. and B. Collins. 1992. Landscape ecology of birds breeding in temperate forest
4 fragments. Pages 443-454 *in* J.M. Hagan III, and D.W. Johnson, editors. Ecology and
5 conservation of neotropical migrant landbirds. Smithsonian Institution Press,
6 Washington, D.C., USA.
- 7 Gardner, R.H., M.G. Turner, R.V. O'Neill, and S. Lavorel. 1991. Simulation of scale-dependent
8 effects of landscape boundaries on species persistence and dispersal. Pages 76-89 *in*
9 M.M. Holland, P.G. Risser, and R.J. Naiman, editors. Ecotones: the role of landscape
10 boundaries in the management and restoration of changing environments. Chapman Hall,
11 New York, USA.
- 12 Gobeil, J.-F. and M.-A. Villard. 2002. Permeability of three boreal forest landscape types to bird
13 movements as determined from experimental translocations. *Oikos* **98**:447-458.
- 14 Goodwin, B.J. and L. Fahrig. 2002. How does landscape structure influence landscape
15 connectivity? *Oikos* **99**:552-570.
- 16 Guenette J.-S. and M.-A. Villard. 2005. Thresholds in forest bird response to habitat alteration as
17 quantitative targets for conservation. *Conservation Biology* **19**:1168-1180.
- 18 Gustafson, E.J. 1998. Quantifying landscape spatial pattern: What is the state of the art?
19 *Ecosystems* **1**:143-156.
- 20 Haddad, N.M. and K.A. Baum. 1999. An experimental test of corridor effects on butterfly
21 densities. *Ecological Applications* **9**:623-633

- 1 Hagan, J.M. and A.L. Meehan. 2002. The effectiveness of stand-level and landscape-level
2 variables for explaining bird occurrence in an industrial forest. *Forest Science* **48**:231-
3 242.
- 4 Haila, Y. 1983. Land birds on northern islands: a sampling metaphor for insular colonization.
5 *Oikos* **41**:334-351.
- 6 Haining, R. 1990. *Spatial data analysis in the social and environmental sciences*. Cambridge
7 University Press, Cambridge, UK.
- 8 Hames, R.S., K.V. Rosenberg, J.D. Lowe, and A.A. Dhondt. 2001. Site reoccupation in
9 fragmented landscapes: testing predictions of metapopulation theory. *Journal of Animal*
10 *Ecology* **70**:182-190.
- 11 Hanley, J.A. and B.J. McNeil. 1982. The meaning and use of the area under a receiver operating
12 characteristic (ROC) curve. *Radiology* **143**:29-36.
- 13 Hannon, S.J. and F.K.A. Schmiegelow. 2002. Corridors may not improve the conservation value
14 of small reserves for most boreal birds. *Ecological Applications* **12**:1457-1468.
- 15 Hanski I. and O. Ovaskainen. 2000. The metapopulation capacity of a fragmented landscape.
16 *Nature* **404**:755-758
- 17 Harris, R.J. and J.M. Reed. 2002. Effects of forest-clearcut edges on a forest-breeding songbird.
18 *Canadian Journal of Zoology* **80**:1026-1037.
- 19 Heikkinen, R.K., M. Luoto, R. Virkkala, and K. Rainio. 2004. Effects of habitat cover, landscape
20 structure and spatial variables on the abundance of birds in an agricultural-forest mosaic.
21 *Journal of Applied Ecology* **41**:824-835.
- 22 Hill, M.F. and H. Caswell. 1999. Habitat fragmentation and extinction thresholds on fractal
23 landscapes. *Ecology Letters* **2**:121-127.

- 1 Hinsley, S.A. 2000. The costs of multiple patch use by birds. *Landscape Ecology* **15**:765-775.
- 2 Holmes, R.T. and T.W. Sherry. 1986. Bird community dynamics in a temperate deciduous forest:
3 Long-term trends at Hubbard Brook. *Ecological Monographs* **56**:201-220.
- 4 Holmes, R.T., T.W. Sherry, P. Marra, and K. Petit. 1992. Multiple brooding and productivity of
5 a neotropical migrant, the Black-throated Blue Warbler (*Dendroica caerulescens*), in an
6 unfragmented temperate forest. *Auk* **109**:321-333.
- 7 Homan, R.N., B.S. Windmiller, and J.M. Reed. 2004. Critical thresholds associated with habitat
8 loss for two vernal pool-breeding amphibians. *Ecological Applications* **14**:1547-1553.
- 9 Hosmer, D.W. and S. Lemeshow. 2000. *Applied Logistic Regression*, 2 edition.
10 John Wiley & Sons, USA.
- 11 Jonsen, I. and P. D. Taylor. 2000. *Calopteryx* damselfly dispersions arising from multiscale
12 responses to landscape structure. *Conservation Ecology* 4: 4. [online] URL:
13 <http://www.consecol.org/vol4/iss2/art4/>
- 14 Kaluzny, S.P., S.C.Vega, T.P. Cardoso, and A.A. Shelly. 1996. *S-Plus Spatial Stats users*
15 manual. Seattle, MathSoft.
- 16 Klute, D., M. Lovallo, W.Tzilkowski. 2002. Autologistic regression modeling of American
17 Woodcock habitat use with spatially dependent data. Pages 335-343 in J.M. Scott, P.J.
18 Heglund, M.L. Morrison, J.B. Haufler, M.G. Raphael, W.A. Wall and F.B. Sampson,
19 editors. *Predicting Species Occurrences: issues of accuracy and scale*. Island Press,
20 Washington, USA.
- 21 Krawchuk, M. A., and P. D. Taylor. 2003. Changing importance of habitat structure across
22 multiple spatial scales for three species of insects. *Oikos* **103**:153-161.

- 1 Larson M.A., F.R. Thompson, J.J. Millspaugh, W.D. Dijak, S.R. Shifley. 2004. Linking
2 population viability, habitat suitability, and landscape simulation models for
3 conservation planning. *Ecological Modelling* **180**:103-118.
- 4 Legendre, P. 1993. Spatial autocorrelation: trouble or new paradigm? *Ecology* **74**:1659-1673.
- 5 Lee, M., L. Fahrig, K. Freemark and D.J. Currie. 2002. Importance of patch scale vs landscape
6 scale on selected forest birds. *Oikos* **96**:110-118.
- 7 Levin, S.A. 1992. The problem of pattern and scale in ecology. *Ecology* **73**:1943-1967.
- 8 Levins, R. 1969. Some demographic and genetic consequences of environmental heterogeneity
9 for biological control. *Bulletin of the Entomological Society of America* **15**:237-240.
- 10 Lichstein, J.W., T.R. Simons, and K.E. Franzreb. 2002a. Landscape effects on breeding songbird
11 abundance in managed forests. *Ecological Applications* **12**:836-857.
- 12 Lichstein, J.W., T.R. Simons, S.A. Shriener and K.E. Franzreb. 2002b. Spatial autocorrelation
13 and autoregressive models in ecology. *Ecological Monographs* **72**:445-463.
- 14 Lindenmayer, D.B., R.B. Cunningham, C.F. Donnelly, H. Nix, and B.D. Lindenmayer. 2002.
15 Effects of forest fragmentation on bird assemblages in a novel landscape context.
16 *Ecological Monographs* **72**:1-18.
- 17 Lorimer, C. 1977. The presettlement forest and natural disturbance cycle of northeastern Maine.
18 *Ecology* **58**:139-148.
- 19 MacArthur, R.H. and E.O. Wilson. 1967. *The Theory of Island Biogeography*. Princeton
20 University Press, Princeton, N.J., USA.
- 21 MacArthur, R.H., J.M. Diamond, and J.R. Karr. 1972. Density compensation in island faunas.
22 *Ecology* **53**:330-342.

- 1 MacFaden, S.W. and D.E. Capen. 2002. Avian habitat relationships at multiple scales in a New
2 England forest. *Forest Science* **48**:243-253.
- 3 MacNally, R. 2000. Regression and model-building in conservation biology, biogeography and
4 ecology: The distinction between – and reconciliation of – ‘predictive’ and ‘explanatory’
5 models. *Biodiversity and Conservation* **9**:655-671.
- 6 Manel, S., H.C. Williams, and S.J. Ormerod. 2001. Evaluating presence-absence models in
7 ecology: the need to account for prevalence. *Journal of Applied Ecology* **38**:921-931.
- 8 Manolis, J.C., D.E. Andersen, and F.J. Cuthbert. 2002. Edge effect on nesting success of ground
9 nesting birds near regenerating clearcuts in a forest-dominated landscape. *Auk* **119**:955-
10 970.
- 11 Marzluff, J.M., J.J. Millspaugh, P. Hurlvitz, and M.S. Handcock. 2004. Relating resources to a
12 probabilistic measure of space use: forest fragments and Stellar’s jays. *Ecology* **85**:1411-
13 1427.
- 14 Matthews, E., R. Payne, M. Rohweder, and S. Murray. 2001. Pilot analysis of global ecosystems:
15 Forest ecosystems. World Resources Institute.
- 16 McComb, W.C. 1999. Synthesis of the Conference. Pages 295-301 in J.A. Rochelle, L.A.
17 Lehmann, and J. Wisniewski, editors. *Forest Fragmentation : wildlife and management*
18 *implications*. Brill Academic Publishing, Leiden, The Netherlands.
- 19 McGarigal, K. and W.C. McComb. 1995. Relationships between landscape structure and
20 breeding birds in the Oregon Coast Range. *Ecological Monographs* **65**: 215-260.
- 21 Mitchell, M.S., R.A. Lancia, and J.A. Gerwin, 2001. Using landscape-level data to predict the
22 distribution of birds on a managed forest: effects of scale. *Ecology* **11**:1692-1708.

- 1 Moilanen, A. and I. Hanski. 1998. Metapopulation dynamics: effects of habitat quality and
2 landscape structure. *Ecology* **79**:2503-2515.
- 3 Mönkkönen, M., P. Helle, G.J. Niemi, and K. Montgomery. 1999. Evolution of heterospecific
4 attraction: using other species as cues in habitat selection. *Evolutionary Ecology* **13**:91-
5 104.
- 6 Mönkkönen, M., and P. Reunanen. 1999. On critical thresholds in landscape connectivity: a
7 management perspective. *Oikos* **84**:302-305.
- 8 Morse, D.H.1994. Blackburnian warbler (*Dendroica fusca*). in A. Poole and F. Gill, editors.
9 Birds of North America, No.102. Academy of. Natural Science and American
10 Ornithological Union, Philadelphia, PA, USA.
- 11 New Brunswick Department of Natural Resources. 1993. New Brunswick Forest Inventory.
12 Fredericton, N.B., Canada.
- 13 Nol, E., C.M. Francis, and D.M. Burke. 2005. Using distance from putative source woodlots to
14 predict occurrence of forest birds in putative sinks. *Conservation Biology* **19**:836-844.
- 15 Norris, D.R. and B.J.M. Stutchbury. 2001. Extraterritorial movements of a forest songbird in a
16 fragmented landscape. *Conservation Biology* **15**:729-736.
- 17 Norton, M.R., S.J. Hannon and F.K.A. Schmiegelow. 2000. Fragments are not islands: patch vs
18 landscape perspectives on songbird presence and abundance in a harvested boreal forest.
19 *Ecography* **23**:209-223.
- 20 Pimm, S.L., G.J. Russell, J.L. Gittleman, , and T.M. Brooks. 1995. The future of biodiversity.
21 *Science* **269**:347-350.
- 22 Pulliam, H.R. and B.J. Danielson. 1991. Sources, sinks and habitat selection: a landscape
23 perspective on population dynamics. *American Naturalist* **137**:S50-S66.

- 1 R Development Core Team. 2004. R: A language and environment for statistical computing. R
2 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-00-3, URL
3 <http://www.R-project.org>.
- 4 Ralph C.J., J. Sauer, and S. Droege (editors).1995. Monitoring Bird Populations by Point
5 Counts. Gen. Tech. Rep. PSW-GTR-149. Pacific Southwest Research Station, Forest
6 Service, U.S. Department of Agriculture, Albany, C.A.
- 7 Reunanen, P., A. Nikula, M. Mönkkönen, E. Hurme and V. Nivala. 2002. Predicting occupancy
8 for the Siberian flying squirrel in old-growth forest patches. *Ecological Applications* **12**:
9 1188-1198.
- 10 Ricketts, T. H. 2001. The matrix matters: effective isolation in fragmented landscapes. *American*
11 *Naturalist* **158**:87-99.
- 12 Robichaud, I., M.-A.Villard, and C.S. Machtans. 2002. Effects of forest regeneration on songbird
13 movements in a managed forest landscape of Alberta, Canada. *Landscape Ecology* **17**:
14 247-262.
- 15 Rodenhouse, N.L., T.S. Sillett, P.J. Doran, and R.T. Holmes. 2003. Multiple density-dependence
16 mechanisms regulate a migratory bird population during the breeding season.
17 *Proceedings of the Royal Society of London Series B* **270**: 2105-2110
- 18 Root, R.B. 1973. Organization of a plant-arthropod association in simple and diverse habitats:
19 the fauna of collards. *Ecological Monographs* **45**:95-120.
- 20 Sallabanks,R., J.R.Walters, and J.A. Collazo. 2000. Breeding bird abundance in bottomland
21 hardwood forests: habitat, edge, and patch size effects. *Condor* **102**:748-758.

- 1 Sauer, J.R., J.E. Hines, and J. Fallon. 2004. The North American Breeding Bird Survey, Results
2 and Analysis 1966 - 2003. Version 2004.1. USGS Patuxent Wildlife Research Center,
3 Laurel, MD, USA.
- 4 Schmiegelow, F.K.A. and M. Mönkkönen. 2002. Habitat loss and fragmentation in dynamic
5 landscapes: avian perspectives from the boreal forest. *Ecological Applications* **12**:375-
6 389.
- 7 Suorsa, P., E. Huhta, A. Jäntti, A. Nikula, H. Helle, M. Kuitunen, V. Koivunen, and H.
8 Hakkarainen. 2005. Thresholds in selection of breeding habitat by the Eurasian
9 treecreeper (*Certhia familiaris*). *Biological Conservation* **121**:443-452.
- 10 Tewksbury, J.J., S.J. Heil, and T.E. Martin. 1998. Breeding productivity does not decline with
11 increasing fragmentation in a western landscape. *Ecological Applications* **79**:2890-2903.
- 12 Tischendorf, L., D.J. Bender, and L. Fahrig. 2003. Evaluation of patch isolation metrics in
13 mosaic landscapes for specialist vs. generalist dispersers. *Landscape Ecology* **18**:41-50.
- 14 Trzcinski M.K., L. Fahrig and G. Merriam. 1999. Independent effects of forest cover and
15 fragmentation on the distribution of forest breeding birds. *Ecological Applications* **9**:586-
16 593.
- 17 Tubelis, D.P., D.B. Lindenmayer, and A. Cowling. 2004. Novel patch-matrix interactions: patch
18 width influences matrix use by birds. *Oikos* **107**:634-644.
- 19 Turner, M. 1989. Landscape ecology: The effect of pattern on process. *Annual Review of*
20 *Ecology Evolution and Systematics* **20**:171-197.
- 21 Van Horne, M.A. and T.M. Donovan. 1994. Ovenbird (*Seiurus aurocapillus*). in A. Poole, F.,
22 Gill, editors. *Birds of North America*, No.88. Academy of. Natural Science and
23 American Ornithological Union, Philadelphia, PA.

- 1 Vega, R.J.H.A., J.H. Rappole, W.J.McShea, and C.A.Haas, 1998. Wood thrush postfledging
2 movements and habitat use in northern Virginia. *Condor* **100**:69-78.
- 3 Venables, W.N. and B.D. Ripley. 2001. *Modern Applied Statistics in Splus*. Springer Verlag,
4 New York, USA.
- 5 Villard, M.-A., G.Merriam, and B.A.Maurer. 1995. Dynamics in subdividing populations of
6 neotropical migratory birds in a fragmented temperate forest. *Ecology* **76**:27-40.
- 7 Villard, M.-A., M.K. Trzcinski, and G. Merriam. 1999. Fragmentation effects on forest birds:
8 relative influence of woodland cover and configuration on landscape occupancy.
9 *Conservation Biology* **13**:774-783.
- 10 Webster, M.S., H.C. Chuang-Dobbs, and R.T. Holmes. 2001. Microsatellite identification of
11 extrapair sires in a socially monogamous warbler. *Behavioural Ecology* **12**:439-446
- 12 Wiegand, T., E. Revilla, and K.A. Moloney. 2005. Effects of habitat loss and fragmentation on
13 population dynamics. *Conservation Biology* **19**:108-121.
- 14 Wiens, J.A. 1994. Habitat fragmentation: island v landscape perspectives on bird conservation.
15 *Ibis* **137**:S97-S104.
- 16 Wiens, J.A., N.C. Stenseth, B. Van Horne, and R.A. Ims. 1993. Ecological mechanisms in
17 landscape ecology. *Oikos* **66**:369-380.
- 18 With, K.A. and A.W. King. 1999. Extinction thresholds for species in fractal landscapes.
19 *Conservation Biology* **13**:314-326.
- 20 Woolfenden, B.E., B.J.M. Stutchbury and E.S. Morton. 2005. Male Acadian flycatchers,
21 *Empidonax virescens*, obtain extrapair fertilizations with distant females. *Animal*
22 *Behaviour* **69**:921-929.

- 1 Young, L., M.G. Betts, and A.W. Diamond. 2005. Blackburnian warbler is a mixed forest
2 specialist. Submitted to *Forest Ecology and Management* **214**:358-372.
- 3 Zabel, C.J., J.R. Dunk, H.B. Stauffer, L.M. Roberts, B.S. Mulder, and A. Wright. 2003. Northern
4 spotted owl habitat models for research and management application in California (USA).
5 *Ecological Applications* **13**:1027-1040.

1 **Tables**

2

3 TABLE 1. Correlation (r) among landscape configuration and composition variables for
 4 blackburnian warbler (bold below diagonal), and ovenbird (italics, above diagonal) ($N=363$).
 5

	HAB300	HAB2000	MATRIX300	MATRIX2000	Patch [†]	Edge ^{††}
HAB300	1.00	<i>0.50</i>	<i>-0.65</i>	<i>-0.34</i>	<i>0.71</i>	<i>0.42</i>
HAB2000	0.64	1.00	<i>-0.31</i>	<i>-0.80</i>	<i>0.55</i>	<i>0.21</i>
MATRIX300	-0.70	-0.44	1.00	<i>0.41</i>	<i>-0.50</i>	<i>-0.49</i>
MATRIX2000	-0.42	-0.84	0.47	1.00	<i>-0.44</i>	<i>-0.31</i>
Patch	0.71	0.57	-0.54	-0.48	1.00	<i>0.37</i>
Edge	0.40	0.31	-0.52	-0.37	0.32	1.00

6

7 *Notes:* HAB = Habitat. This was calculated as the summed estimated probability of
 8 occurrence for both species (\hat{p}) for all 30 m² pixels within a 300 or 2000 m radius. MATRIX =
 9 Amount of non-habitat ($\hat{p} < 0.05$) within a 300 or 2000 m radius. See *Methods* for details.

10 Numbers beside variable names indicate spatial extent of variables (the radius of a circle centered
 11 on each sample point).

12 [†]Patch size

13 ^{††}Distance to hard edge

14

1 TABLE 2. Coefficients (β) and lower (LCI) and upper (UCI) 95% confidence intervals for Blackburnian Warbler Generalized Linear
 2 Mixed Models with the lowest AIC in 2002, 2003 and for reoccurrence.

Variable	2002			2003			Reoccurrence		
	β	LCI	UCI	β	LCI	UCI	β	LCI	UCI
Intercept	0.672			-0.266			1.065		
Trees>30	0.205	0.051	0.359	0.286	0.127	0.445	0.303	0.129	0.478
Canopy cover (%)	0.665	-0.250	1.580	-	-	-	-	-	-
HWD>10	-0.242	-0.488	0.004	-0.142	-0.396	0.111	-0.494	-0.812	-0.176
SWD>20	-0.177	-0.263	-0.090	-0.123	-0.215	-0.032	-0.235	-0.357	-0.112
MATRIX300	-0.005	-0.009	-0.001	-0.005	-0.010	-0.001	-0.005	-0.011	0.000
MATRIX2000	-0.0001	-0.0003	0.0000	-	-	-	-0.0001	-0.0003	0.0000
HWD>10 * SWD>20	0.052	0.031	0.074	0.041	0.019	0.063	0.067	0.038	0.096

3
 4 *Notes:* Trees>30 = number of trees >30 cm dbh/ ha, HWD>10 = number of hardwood trees >10 cm dbh/ ha, SWD>20 = number
 5 of softwood trees >20 cm dbh within a 50 m radius. For other abbreviations see Table 1.

1 TABLE 3. Akaike weights (w_i), evidence ratios (Evidence), and number of parameters (K) from AIC-based model selection for
 2 blackburnian warbler occurrence in 2002, 2003 and for reoccurrence.

3

Hypothesis	2002			2003			Reoccurrence		
	w_i	Evidence	K	w_i	Evidence	K	w_i	Evidence	K
Random sample	0.00	169.7	6	0.03	11.9	5	0.01	40.0	5
Composition	0.22	1.0	8	0.39	1.0	6	0.30	1.0	7
Fragmentation	0.03	6.8	8	0.06	7.0	8	0.05	6.2	7
Non-linear fragmentation	0.03	7.0	9	0.01	38.1	8	0.02	16.8	8

4

5 *Notes:* Only models with the lowest AIC for each landscape hypothesis are shown. All models within a year and for reoccurrence
 6 contain the same local-scale variables. Weights of models with the greatest support given the data are highlighted.

1

2 TABLE 4. Coefficients (β) and lower (LCI) and upper (UCI) 95% confidence intervals for ovenbird Generalized Linear Mixed Models
 3 with the lowest AIC in 2002, 2003 and for reoccurrence.

Variable	2002			2003			Reoccurrence		
	β	LCI	UCI	β	LCI	UCI	β	LCI	UCI
Intercept	-7.338			-4.026			-5.927		
Hardwood BA	0.035	0.008	0.062	0.030	0.008	0.052	0.029	0.007	0.051
Litter (%)	1.561	0.497	2.625	-	-	-	0.610	0.065	1.286
Canopy cover (%)	0.943	0.195	1.691	0.851	-0.052	1.754	1.061	0.116	2.006
OVEN300	0.011	0.004	0.018	-	-	-	-	-	-
OVEN2000	0.0007	0.0003	0.0011	0.0004	0.0001	0.0007	0.0006	0.0002	0.0009
Patch size (ha) [†]	1.745	0.644	2.846	1.156	0.358	1.954	1.620	0.684	2.557
OVEN2000 * Patch size	-0.0005	-0.0003	-0.0002	-0.0002	-0.0003	-0.00005	-0.0003	-0.0004	-0.0001

4

5 *Notes:* Hardwood BA = hardwood basal area/ ha (m²). For other variable abbreviations see Table 1.

6 [†]Log transformed.

1

2 TABLE 5. Akaike weights (w_i), evidence ratios (Evidence), and number of parameters (K)

3 from AIC-based model selection for ovenbird occurrence in 2002, 2003 and for reoccurrence.

4

Hypothesis	2002			2003			Reoccurrence		
	w_i	Evidence	K	w_i	Evidence	K	w_i	Evidence	K
Random sample	0.00	$2.89 \cdot 10^4$	4	0.03	6.66	3	0.02	30.72	4
Composition	0.01	66.06	5	0.18	1.03	4	0.02	26.03	5
Fragmentation	0.00	274.55	7	0.03	6.16	6	0.01	75.04	6
Non-linear fragmentation	0.89	1.00	8	0.20	1.00	6	0.55	1.00	7

5

6 *Notes:* Only models with the lowest AIC for each landscape hypothesis are shown.

7 All models within a year and for reoccurrence contain the same local-scale variables.

8 Weights of models with the greatest support given the data are highlighted.


1 **Figure Captions**

2 FIG. 1. Variance partitioning results showing the proportion of independently explained
3 variance by local-scale, landscape composition, landscape fragmentation, and non-linear
4 fragmentation (Non-linear frag) variables for blackburnian warbler (A) and ovenbird (B) in
5 2002, 2003 and for reoccurrence.


6

7 FIG. 2. Interaction between effect of patch size and landscape extent habitat amount (%
8 within a 2000 m radius) on the raw proportion (\pm SE) of ovenbirds observed in 2002. For
9 graphical purposes, proportions were calculated from the ratio of presences to absences
10 within five categories: 0-5 ha, 6-20 ha, 21-50 ha, 51-100 ha, >100 ha and three habitat
11 amount categories (above). SE was calculated as $\sqrt{(p * q) / N}$ where p = proportion of
12 presences in a category, and q = the proportion of absences. Proportions do not control for
13 local site variation as was conducted in statistical models.

14



1
2
3



1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

APPENDIX A

A table showing local and landscape variables used to predict build models to predict the occurrence and reoccurrence of blackburnian warbler and ovenbird are available in ESA’s Electronic Data Archive: *Ecological Archives*.

APPENDIX B

AIC model rankings, weights (w_i) and evidence ratios (ER) of blackburnian warbler models for 2002 are available in ESA’s Electronic Data Archive: *Ecological Archives*.

APPENDIX C

AIC model rankings, weights (w_i) and evidence ratios (ER) of blackburnian warbler models for 2003 are available in ESA’s Electronic Data Archive: *Ecological Archives*.

APPENDIX D

AIC model rankings, weights (w_i) and evidence ratios (ER) of blackburnian warbler models for reoccurrence are available in ESA’s Electronic Data Archive: *Ecological Archives*.

APPENDIX E

AIC model rankings, weights (w_i) and evidence ratios (ER) of ovenbird models for 2002 are available in ESA’s Electronic Data Archive: *Ecological Archives*.

1

APPENDIX F

2 AIC model rankings, weights (w_i) and evidence ratios (ER) of ovenbird models

3 for 2003 are available in ESA's Electronic Data Archive: *Ecological Archives*.

4 .

5

APPENDIX G

6 AIC model rankings, weights (w_i) and evidence ratios (ER) of ovenbird

7 reoccurrence models are available in ESA's Electronic Data Archive: *Ecological*

8 *Archives*.

9



Partners in Sustainability/Partenaires pour durabilité

Atlantic Society of Fish and Wildlife Biologists
Canadian Institute of Forestry
Canadian Forest Service
City of Moncton
Conservation Council of New Brunswick
Eel Ground First Nation
Elgin Eco Association
Environment Canada
Fawcett Lumber Company
Fisheries and Oceans Canada
Fundy Environmental Action Group
Fundy National Park
Greater Fundy Ecosystem Research Group
Indian and Northern Affairs Canada
INFOR Inc.
J.D. Irving, Limited
K.C. Irving Chair in Sustainable Development, Université de Moncton
Maritime College of Forest Technology
NB Department of Environment and Local Government
NB Department of Natural Resources and Energy
NB Federation of Naturalists
NB Federation of Woodlot Owners
NB Premier's Round Table on the Environment & Economy
NB School District 2
NB School District 6
Nova Forest Alliance
Petitcodiac Sportsman's Club
Red Bank First Nation
Remsoft Inc.
Southern New Brunswick Wood Cooperative Limited
Sussex & District Chamber of Commerce
Sussex Fish and Game Association
Town of Sussex
Université de Moncton
University of New Brunswick Fredericton Faculty of Forestry and Environmental
Management
University of New Brunswick Saint John
Village of Petitcodiac
Washademoak Environmentalists